Table 15: Alcohol Related Risk Behaviors, Montana Adults, 2010 (with 95% confidence intervals)

	Binge Drinking				Heavy Drinking				Any Drinking and Driving			
	Past 30 Days † 95% CI				Past 30 Days ‡ 95% CI				Past 30 Days § 95% CI			
	Wt.%	95% LL		UnWt. N	Wt.%	95% LL		UnWt. N	Wt.%	95% LL		UnWt. N
All Adults	17.0	15.4	18.6	956	5.3	4.6	6.1	395	4.8	3.8	6.0	170
	17.0	15.4	10.0	930	5.5	4.0	0.1	393	4.0	3.0	0.0	170
Sex:												
Male	22.6	20.1	25.4	569	5.7	4.6	7.0	189	6.3	4.8	8.2	123
Female	11.4	9.8	13.3	387	4.9	4.1	5.9	206	3.0	2.0	4.5	47
Age:												
18 - 24	22.9	15.6	32.2	56	2.6	1.4	5.0	14	7.3	3.5	14.7	11
25 - 34	25.9	21.0	31.5	149	6.0	3.8	9.5	30	5.9	3.3	10.3	23
35 - 44	23.3	19.7	27.3	187	5.9	4.1	8.4	51	5.7	3.4	9.3	29
45 - 54	18.4	15.9	21.2	263	7.0	5.5	8.9	108	5.5	3.9	7.8	51
55 - 64	11.6	9.9	13.6	202	6.2	4.9	7.7	102	3.2	2.2	4.8	38
65+	3.9	3.0	5.0	96	3.3	2.6	4.3	89	1.2	0.7	2.2	17
Education:												
<high school<="" td=""><td>14.8</td><td>10.3</td><td>20.8</td><td>66</td><td>3.7</td><td>2.0</td><td>6.7</td><td>22</td><td>3.8</td><td>1.8</td><td>7.8</td><td>10</td></high>	14.8	10.3	20.8	66	3.7	2.0	6.7	22	3.8	1.8	7.8	10
High School	17.5	14.7	20.6	300	4.8	3.6	6.3	116	4.7	3.1	7.1	40
Some College	19.6	16.4	23.1	312	6.2	4.7	8.1	116	7.4	5.1	10.7	62
College Degree	14.6	12.5	17.1	278	5.4	4.3	6.6	141	2.9	2.0	4.1	58
, ,				-								
Income:	40.5	40.0	40.0	400	4.0	0.0	7.0	47	0.4	0.7	40.0	47
<\$15,000	13.5	10.0	18.0	106	4.9	3.3	7.2	47 55	8.4	3.7	18.0	17
\$15,000 - \$24,999	15.2	11.5	19.8	139	4.4	2.9	6.6	55	4.0	2.0	7.5	20
\$25,000 - \$49,999	17.0	14.4	20.0	287	4.9	3.9	6.1	123	4.8	3.2	6.9	52
\$50,000 - \$74.999	16.1	13.0	19.7	160	6.8	4.7	9.8	65 86	4.8	2.8	8.1	30 49
\$75,000 +	23.4	19.5	27.8	226	6.4	4.8	8.6	86	5.4	3.6	8.0	49
Race/Ethnicity:												
White, non-Hispanic	16.3	14.7	18.0	783	5.3	4.6	6.2	346	4.7	3.7	6.0	140
AI/AN*	23.2	17.3	30.2	118	4.5	2.3	8.4	29	9.8	5.3	17.6	25
Other or Hispanic**	NSD [∆]			50	5.9	2.8	11.7	19	2.9	0.7	10.9	4
Disability:												
Disability	11.1	9.0	13.8	211	4.3	3.4	5.5	110	3.4	2.2	5.1	41
No Disability	19.1	17.2		741	5.6	4.7	6.7	284	5.2	4.0	6.6	129
Ĭ												
Region:	16 F	12.2	20.4	107	E 6	2.0	0.0	44	0.6	ΕO	115	25
1- Eastern MT	16.5	13.2		127	5.6	3.8	8.2	41	8.6	5.0	14.5	25
2- N Central MT 3- S Central MT	20.5 14.6	17.0 11.2	24.5 18.8	243 105	4.5 4.1	3.1 2.5	6.3 6.6	74 36	5.2 6.3	3.3 3.4	8.1 11.3	43 20
4- Southwest MT	15.5	12.8	18.7	182	4.1 4.7	2.5 3.6	6.2	36 78	3.8	3.4 2.4	5.9	30
5- Northwest MT	17.6	14.7	20.9	287	4.7 6.7	5.3	8.4	78 159	3.8	2.4	5.9	50 50
MMSA-Billings	14.8	10.7	20.9	51	4.1	2.2	7.3	18	6.0	2.8	12.5	8
MMSA-Helena	17.3	13.1	22.5	80	5.2	3.2	8.2	31	7.8	4.2	14.0	16
MMSA-Helena MMSA-Kalispell	16.6	12.8		81	6.5	3.2 4.5	9.2	43	3.1	1.5	6.3	10
wiwion-Nalispeli	10.0	12.0	۷۱.۵	01	0.5	٦.٠	٥.۷	70	J. 1	1.5	0.0	10

[†] Binge drinking is defined as having five or more alcoholic drinks on one occasion for men, and four or more alcoholic drinks on one occasion for women. Total Sample Size: 7,169, Weighted Prevalence Estimate: 127,000.

[‡] Heavy drinking is defined as the consumption of more than two alcoholic drinks per day for men or more than one alcoholic drink per day for women. Total Sample Size: 7,124, Weighted Prevalence Estimate: 39,000.

[§] Among adults that had at least one alcoholic drink in the past 30 days. During the past 30 days, how many times have you driven when you've had perhaps too much to drink? Total Sample Size: 3,772, Weighted Prevalence Estimate: 20,000.

^{*} American Indian or Alaska Native only.

^{**} All other non-White (including multiracial) or Hispanic.

 $[\]Delta$ Not Sufficient Data to report a reliable estimate.